

Mould Growth and Aflatoxin Accumulation Affected by Newly Synthesized Derivative of Coumarine Treatment of Maize Hybrid

Lejla DURAKOVIĆ¹ (✉), Mihaela BLAŽINKOV², Branka ŠEOL³, Alma TUDIĆ⁵,
Frane DELAŠ¹, Marijan BOŠNJAK¹, Sanja SIKORA², Andrea SKELIN²,
Katarina HUIĆ-BABIĆ², Zijad DURAKOVIĆ⁴

Summary

The effect of newly synthesized derivative of 11-hydroxy-7-imino-7H-7a, 12-diazabenzofuran-6-one, on growth and aflatoxin B₁ and G₁ (AFB₁ and AFG₁) accumulation by toxigenic mould *Aspergillus flavus* ATCC 26949 was studied on a solid substrate (maize grains) to determine the possible use of this compound as a mean of controlling aflatoxin accumulation. Experiments were carried out in a stationary culture at temperature of 28°C during 21 days. The growth of mould was monitored by measuring the analysis of chitin as glucosamine, as a criterion, and concentration of AFB₁ and AFG₁ was measured by HPLC method using Hewlett-Packard instrument with fluorescence detector. Concentration of investigated coumarine of 0.05 mmol x g⁻¹ stimulated mould growth and aflatoxin accumulation, but concentration of 0.2 mmol x g⁻¹ or higher produced an inhibitory effect. In the presence of 0.2 and 0.5 mmol x g⁻¹ of this compound, mould growth was decreased by 22% and 65%, respectively. Concentration of AFB₁ in these investigations was reduced by 30% and 90%, and concentration of AFG₁ was reduced almost completely in respect to values obtained in control experiments. In experiments with 2.0 mmol x g⁻¹ of this inhibitor no synthesis of both investigated aflatoxins was found in the sample although small increment of biomass was detected.

Key words

moulds, chitin, aflatoxins, antifungal agents, coumarine

¹ University of Zagreb, Faculty of Food Technology and Biotechnology, Department of Biochemical Engineering, Pierottieva 6, 10000 Zagreb, Croatia

✉ e-mail: Lejla.Durakovic@pbf.hr

² University of Zagreb, Faculty of Agriculture, Department of Microbiology, Svetošimunska 25, 10000 Zagreb, Croatia

³ University of Zagreb, Veterinary Faculty, Department for Microbiology and Infectious Disease with Clinic, Heinzelova 55, 10000 Zagreb, Croatia

⁴ Institute for Anthropology, Department for Medical Anthropology and Epidemiology, Gajeva 32, 10000 Zagreb, Croatia

⁵ Croatian Waters, Sector of Development and Water Economy, Vukovarska 220, 10000 Zagreb, Croatia

Received: October 29, 2010 | Accepted: December 5, 2010

ACKNOWLEDGEMENTS

The authors would like to thank Professor E.B. Lillehoy, United States Department of Agriculture, Peoria, Illinois for supplying the strain of *Aspergillus flavus* used in this work. We also grateful to Professor M. Terzić for analyses of aflatoxins in maize grains and for information, and Professor V. Rapić for providing of investigated coumarine. Finally, the authors gratefully acknowledge gentlewoman Brigitta Duraković for language correcting the manuscript. The investigations were supported by a grant No. 058-0582184-0432 from Croatian Ministry of Science and Technology.

Introduction

Foods and feeds are frequently contaminated with moulds during harvesting, storage and handling before reaching the consumer.

Many of these ubiquitous moulds have been shown to produce toxins. The presence of known toxin-producing moulds does not necessarily indicate the presence of mycotoxins, nor does the lack of a mould indicate that the sample has not been contaminated with a mycotoxin. Some moulds produce toxins under restricted conditions and only particular strains of some species produce toxins (Pittet, 1998).

Among the fungal toxins associated with human and animal food materials are included several derivatives of coumarin that evoke varied pharmacological and physiological responses in the animal body. Aflatoxins are toxic, highly carcinogenic secondary metabolites produced by the fungal genera *Aspergillus*, *Penicillium* and *Rhizopus* (Goldblatt, 1969; Brown *et al.*, 1999; Dorner *et al.*, 1999; Wu, 2004; Binder, 2007). AFB₁ has been associated with certain mycotoxicoses including turkey "X" disease and chronic exposure to low levels of this toxin pose a serious health and economic hazard (Diener *et al.*, 1987; Karlowsky, 1999; Thompson and Henke, 2000; CAST Report, 2003).

To prevent aflatoxin formation in agricultural commodities it is first necessary to prevent the growth aflatoxin producing fungi (Windham *et al.*, 1999; Wright *et al.*, 2000; Schatzmayr *et al.*, 2006). This can be achieved by at least three means: control of the environment, use of chemical antifungal agents and utilization of natural resistance in agricultural commodities (Duraković *et al.*, 1985; Sušac *et al.*, 1989; Samarajeva *et al.*, 1990; Praxton, 1991; Gourama and Bullerman, 1995; Cho and Kang, 2000; Wright *et al.*, 2000; Delaš *et al.*, 2008a, 2008b, 2008c; Duraković *et al.*, 2008). During the past 40 years there has been increasing interest in the identification of naturally occurring or newly synthesized chemical compounds that inhibit growth and/or aflatoxin production by aflatoxigenic species (Hitokoto *et al.*, 1980; Duraković *et al.*, 1985; Sušac *et al.*, 1989; Duraković *et al.*, 2006; Duraković, 2007; Duraković *et al.*, 2007; Duraković *et al.*, 2008; Duraković *et al.*, 2010).

Preliminary studies in Laboratory of Microbiology, Faculty of Food Technology and Biotechnology, Zagreb, had indicated the efficacy of such derivatives of coumarine as the inhibitors of the growth aflatoxin-producing fungi (Sušac *et al.*, 1989; Duraković *et al.*, 2006; Delaš *et al.*, 2008a, 2008b, 2008c). This was the basis of our study, which aimed at evaluating the mould growth and aflatoxin accumulation during the growth of *A. flavus* ATCC 26949 on maize in the presence of newly synthesized derivative of coumarine (Duraković, 2007; Duraković *et al.*, 2007; Delaš *et al.*, 2008a).

Materials and methods

Culture and inoculum. *Aspergillus flavus* ATCC 26949 strain was obtained from the USDA Fermentation Laboratory Northern Regional Research Center, Peoria, IL. The cultures were maintained on slants of potato dextrose agar (PDA) stored at 4°C. Before each experiment, the mould was transferred to another PDA slant and was incubated at ambient temperature.

The inoculum for all experiments was prepared by growing *A. flavus* on PDA in 100 x 15 mm Petri dishes for seven days at 28°C until it was well sporulated. Contents of two Petri dishes, including agar and fungus, were homogenized in 50 mL sterile water with a Potter homogenizer for 1-2 minutes. The culture was shaken vigorously for 1-2 minutes, and spore number was determined using a Thoma counting chamber. The spore suspension was diluted to 5 x 10⁸ conidia x mL⁻¹.

Assay of antifungal activity. The antifungal activity of investigated coumarine was assayed using two techniques as follows:

- Standard assay:** 1 mL of the spore suspension of *A. flavus* ATCC 26949 was used to seed the Petri dishes, prepared by pouring 20 mL of PDA. Agar medium was poured into dishes, and the content of the dish mixed well. After allowing the agar to solidify, dishes were prepared using a sterile glass tube and pipette bulb. The various concentrations of investigated coumarine were then added to the dishes in appropriate quantities in triplicate. In each case, a separate dish with pure solvent was employed as a control. The diameter of zone inhibition was measured after incubation of 48 hours at temperature of 25°C.
- Assay in flask cultures:** The studies of aflatoxin accumulation by *A. flavus* were carried out in duplicate in Erlenmeyer flasks contained 50 g of maize grains. The water content in the substrate was adjusted by adding appropriate amounts of distilled water to the samples to quantity of 40%. Substrates were autoclaved and then inoculated with 1 mL of the spore suspension containing 5 x 10⁸ conidia x mL⁻¹ of *A. flavus*. On the basis of preliminary information obtained from the dishes assay, appropriate quantities of various concentrations of coumarine were added in triplicate to the flasks containing maize grains and spores. The flasks were kept for 24 hours with thorough agitation so that the coumarine could be absorbed by grains. Flasks not containing coumarine served as a control. Flasks were incubated at 25°C for 15 days. A method of measuring the degree of fungal invasion on corn was analysis of chitin, measured as glucosamine (Donald and Mirocha, 1977).

Substrate for mould growth and aflatoxin accumulation. The basal substrate used in this study was whole maize grains (ZG-SK 502A hybrid). Fifty grams of grains were distributed in each 300 mL Erlenmeyer flasks. Sterile distilled water (about 22 mL x 50 g⁻¹) was added to achieve moisture content of 40%. The investigated coumarine was synthesized in the Laboratory of Organic Chemistry of Faculty of Food Technology and Biotechnology in Zagreb, and was dissolved in dimethylsulfoxide (DMSO) + water (1 : 1 w/w) at 1.0 mol x L⁻¹ concentration. Its structure is shown in Figure 1.

A method for the synthesis of this compound is described by Govori *et al.*, 2002 and Chen *et al.*, 2010. The required amounts of this solution to give 0.05; 0.2; 0.5; 1.0 and 2.0 mmol x g⁻¹ of broth were pipetted into test Erlenmeyer flasks. Control test flasks and duplicate test flasks containing the various concentrations of coumarine were inoculated with 1.0 mL of the spore suspension. All the flasks were incubated at 28°C for 21 days.

Measurement of fungal growth. Every seven days control flask and duplicate test flasks were taken out of the incubator as

samples for the determination of the amount of biomass and the AFB₁ and AFG₁ accumulation. The substrates were autoclaved at 121°C for 30 minutes before analysis to kill the spores and vegetative mycelia. The growth of *A. flavus* was monitored by using the analysis of chitin measured as glucosamine, as a criterion (Donald and Mirocha, 1977). Chitin, a polymer of N-acetyl-D-glucosamine, is a constituent of the cell walls of most fungi and can be used as a measure of total fungal growth, since little or no chitin-like materials occur in sound cereal grains. In the analytical devices, the polymer is not measured directly but rather is hydrolysed to glucosamine, deaminated to its corresponding aldehyde and measured spectrophotometrically. The chitin content is estimated from the standard curve of glucosamine-HCl read at 650 nm.

On the basis of the data obtained, the calibration curve was made, from which was, according to the chitin content, directly determined amount of biomass dry weight (Duraković, 1981).

The percentage inhibition of mould growth was determined according to the formula:

$$\% \text{ inhibition} = \left(1 - \frac{m_1}{m_2}\right) \times 100$$

m_1 = biomass dry weight in flask containing inhibitor (mg x g⁻¹)

m_2 = biomass dry weight in control flask (mg x g⁻¹)

Aflatoxin assay. The aflatoxins were extracted from samples with chloroform as described by AOAC, 2005. The chloroform was evaporated under nitrogen gas, the samples were dissolved in methanol and filtered through 0.25 µm organic solvent filter (Millipore Co., Bedford, Mass., USA) prior High Performance Liquid Chromatography (HPLC).

To confirm the identity of aflatoxins Thin Layer Chromatography (TLC) was used. Silica gel precoated plates (250 µm thick; Merck, Darmstadt, Germany) were activated at 115°C for one hour before use. TLC plates were developed in chloroform : acetone : petroleum ether (33 : 6 : 1) (modified Duraković *et al.*, 1985), where after they were examined by comparing the R_f values of a known standards to the unknown samples under UV light at 365 nm. Aflatoxin primary standards to check the linearity were provided by Carl Roth (Karlsruhe, Germany). Six milliliter vial of methanol solution with the following specification: 250 ng x mL⁻¹ of AFB₁ and 250 ng x mL⁻¹ of AFG₁. Working standard solutions of 0.25 and 0.125 ng x mL⁻¹ of each aflatoxin (AFB₁ and AFG₁) were prepared by diluting the primary standard solution with methanol-acetonitrile (1 : 1 v/v).

HPLC analysis was performed by using a Hewlett-Packard 1050 Liquid Chromatography (pump and injection system), (Walborn, Germany) with JASCO FP-920 fluorescence detector (Co. Ltd., Japan) and HP integrator 3395. The HPLC column was C₁₈ Nova-Pak (4.6 x 250 mm) with particle size of 5 µm (Waters, Millipore; Millford, MA). Detection of aflatoxins was carried out at λ_{ex} 360 nm and λ_{em} 420 nm. For aflatoxins the fluorescence detector was set at 360 and 420 nm and the mobile phase consisted of a mixture of deionised water-acetonitrile-methanol (60 : 25 : 15 v/v). The flow rate was 1 mL x min⁻¹ for each mobile phase and the injected volume of working standards was 50 µL.




Figure 1. Structure of investigated coumarine (Sušac *et al.*, 1989; Govori *et al.*, 2002; Duraković *et al.*, 2006; Duraković, 2007).

The change of aflatoxin mass fraction (%) is determined according to the formula:

$$\text{concentration change} = \left(1 - \frac{Y_1}{Y_2}\right) \times 100$$

Y_1 = mass fraction of aflatoxin in substrate containing inhibitor (µg x g⁻¹)

Y_2 = mass fraction of aflatoxin in control substrate (µg x g⁻¹)

Results and discussion

The influence of investigated coumarine on the growth of the mould *A. flavus* ATCC 26949 and aflatoxin accumulation are indicated in Table 1 and Figure 2.

Mould growth and AFB₁ and AFG₁ accumulation were observed for 21 days at temperature of 28°C after inoculation of *A. flavus* on whole maize grains. According to previous findings (Gourama and Bullerman, 1995; Wicklow *et al.*, 1998; Weinderbörner, 2001) the optimal temperature for the biosynthesis of AFB₁ is 28-32°C, whereas for AFG₁ it is 24-28°C.

The increased interest in biopreservation of food system has recently led to the development of new synthesized antimicrobial compounds having different origins. A variety of chemical compounds to prevent spoilage by moulds and biosynthesis of mycotoxins have been investigated. These include herbs, spices, essential oils, pesticides, microbial metabolites (bacteriocins and organic acids) and/or by adding fungal inhibitors such as propionic acid, newly synthesized derivatives of dehydroacetic acid, tetraketone and coumarine (Sušac *et al.*, 1989; Samarajeva *et al.*, 1990; Wilkins and Board, 1991; Duraković *et al.*, 1994; Duraković *et al.*, 1995; Basappa and Shantha, 1996; Georgopapadaku, 1998; Duraković *et al.*, 1999; Cho and Kang, 2000; Lavermicocca *et al.*, 2003; CODEX: CAC/RCP 51-2003, 2005; Duraković *et al.*, 2004; Duraković *et al.*, 2006; Duraković, 2007; Delaš *et al.*, 2008a; Parameshwarappa *et al.*, 2009).

Thus Bullerman *et al.* (1977) investigated the antifungal properties of cinnamon in media and raisin bread. They reported the spice to inhibit the growth of *Aspergillus parasiticus* and aflatoxin production; toxin production being inhibited to a greater extent than mycelial growth. Research by Sušac *et al.*

Table 1. Effect of investigated coumarine on growth and aflatoxin accumulation by *Aspergillus flavus* ATCC 26949

| Coumarine (mmol x g ⁻¹) | Incubation time (days) | Biomass dry weight (mg x g ⁻¹) | Aflatoxin (µg x g ⁻¹) | | %Inhibition ^{a100} | | |
|----------------------------------------|---------------------------|-----------------------------------------------|-----------------------------------|----------------|-------------------------------------------------------------|----------------|----------------|
| | | | B ₁ | G ₁ | Growth | Aflatoxin | |
| | | | | | | B ₁ | G ₁ |
| 0 | 7 | 27.80 | 640.0 | 65.0 | - | - | - |
| | 14 | 30.0 | 1560 | 105.0 | - | - | - |
| | 21 | 47.50 | 1000 | 185.0 | - | - | - |
| 0.05 | 7 | 28.30 | 720.0 | 80.0 | Mould growth and accumulation of aflatoxins were stimulated | | |
| | 14 | 34.0 | 1850 | 130.0 | | | |
| | 21 | 52.0 | 1050 | 190.0 | | | |
| 1.0 | 7 | 9.60 | c ₋ | c ₋ | 65.0 | c ₋ | c ₋ |
| | 14 | 12.40 | 170.0 | c ₋ | 60.0 | 90.0 | c ₋ |
| | 21 | 14.20 | 65.0 | c ₋ | 70.0 | 95.0 | c ₋ |
| 2.0 | 7 | b ₋ | c ₋ | c ₋ | c ₋ | c ₋ | c ₋ |
| | 14 | b ₋ | c ₋ | c ₋ | c ₋ | c ₋ | c ₋ |
| | 21 | 5.70 | c ₋ | c ₋ | 99.88 | c ₋ | c ₋ |

a₁₀₀ - (test culture/control culture x 100); b₋ - No detectable growth; c₋ - No detectable aflatoxin

(1989) and Duraković *et al.* (2006., 2007) on *A. parasiticus* NRRL 2999 and *A. flavus* ATCC 26949 in maize shows that some new synthesized derivatives of coumarine possess an effective antifungal and antimycotoxigenic properties. Our previous results demonstrate clearly that mould growth and aflatoxin accumulation were partially stimulated by amounts of investigated tetra-ketone and coumarine derivatives near the minimal inhibitory concentrations (Duraković *et al.*, 1995, 2006).

Table 1 and Figure 2 show the results of growth of mould *Aspergillus flavus* ATCC 26949 on maize grains at incubation temperature of 28°C in the presence of different concentrations of investigated coumarine. Duraković *et al.* (2006) investigated the antifungal and antiaflatoxigenic properties of some newly synthesized derivatives of coumarine during the growth of mould *Aspergillus parasiticus* NRRL 2999 in YES medium.

Results obtained in our study confirmed that one of selected and investigated coumarines is an effective inhibitor of synthesis of biomass and accumulation of AFB₁ and AFG₁ during the growth of mould *Aspergillus flavus* ATCC 26949 on maize grains.

In control experiments the greatest amount of biomass obtained after 21 days during the growth of mould culture was 47.50 mg x g⁻¹ of substrate. The concentration of AFB₁ reached peak values after 14 days and AFG₁ after 21 days. Maximum concentrations of investigated toxins amounted to: 1560 µg AFB₁ x g⁻¹ mycelium dry weight, and 185.0 µg AFG₁ x g⁻¹ mycelium dry weight (Table 1).

The effect of increasing concentration of investigated inhibitor on the growth of mould *A. flavus* and aflatoxin accumulation were evaluated. Concentration of coumarine of 0.05 µg x g⁻¹ of maize grains produced no inhibitory effect, even stimu-




Figure 2. (a, b and c) Comparative presentation of biomass and synthesis of aflatoxins during the growth of mould *Aspergillus flavus* ATCC 26949 in: a) control; b) samples with 0.2 mmol x g⁻¹ and c) samples with 0.5 mmol x g⁻¹ of investigated coumarine. Cultures were cultivated on maize grains at 28 °C for 21 days, and each point represents the average of two samples.

lated mould growth and aflatoxin accumulation (Table 1). After 21 days of incubation biomass dry weight was higher ca 5% as compared in control, and AFB₁ and AFG₁ were higher 12% and 18.5%, respectively. At a concentration of 0.2 mmol x g⁻¹ in 50 g of maize grains, the growth of the mould was reduced by 22% as compared with control and concentration of aflatoxin B₁ was reduced by 30%. After 21 days of cultivation no synthesis of AFG₁ could have been detected (Figure 2b). The biomass content in experiments with 0.5 mmol x g⁻¹ of investigated coumarine was lessened. The highest values obtained after 21 days were reduced by 65% as compared in control (Figure 2c). In these experiments the concentration of AFB₁ after 14 days was considerably lower, and no synthesis of AFG₁ could have been detected. The highest concentration of AFB₁ was 54.0 µg x g⁻¹ of mycelium dry weight, i.e. about 90% lower than in experiments with 0.2 mmol x g⁻¹ of this compound.

In experiments with 1.0 mmol x g⁻¹ of coumarine no synthesis of both aflatoxins were found in the samples after seven days of cultivation although biomass content was 9.6 mg x g⁻¹ of substrate. After 14 and 21 days the biomass content was 12.4 and 14.2 mg x g⁻¹ of substrate, respectively. Concentration of AFB₁ was 170.0 and 65.0 µg x g⁻¹ of substrate and no synthesis of AFG₁ could have been detected (Table 1).

In experiments with 2.0 mmol x g⁻¹ of this compound there was no synthesis neither biomass nor both aflatoxins after seven and 14 days of cultivation. After 21 days of cultivation small increment of biomass was detected, and no accumulation of both investigated aflatoxins (Table 1)

The data obtained in these investigations are standing in good accordance with the findings of several authors (Sušac *et al.*, 1989; Duraković *et al.*, 2006; Duraković, 2007; Parameshwarappa *et al.*, 2009; Roussaki *et al.*, 2010; Tamura *et al.*, 2010) who have shown that some coumarines strongly inhibit growth of toxigenous moulds and biosynthesis of mycotoxins as aflatoxins.

Our results confirmed that selected investigated coumarine is an effective inhibitor of synthesis of biomass and aflatoxin accumulation in maize contaminated with toxigenous mould *Aspergillus flavus* ATCC 26949.

Conclusions

The purpose of this study was to examine the effectiveness of newly synthesized derivative of coumarine for the control of growth of toxigenous mould *Aspergillus flavus* ATCC 26949 and aflatoxin accumulation on maize grain ZG-SK 502A hybrid. Using different concentrations we found that application of investigated coumarine prior to storage may be a potential means of preventing the growth and aflatoxin accumulation by *Aspergillus flavus* ATCC 26949 in the maize grain. Studies are now being conducted to evaluate the coumarine as antifungal and antimycotoxigenic agent.

These data indicate that on maize grain and under certain conditions, investigated coumarine may provide some antifungal and antimycotoxigenic benefit, and thus health protection against possible aflatoxin accumulation. Further investigations on maize grain hybrids might be useful to study the effectiveness of coumarine on growth and aflatoxin accumulation during

the growth of toxigenous mould *Aspergillus flavus* in controlling parameters of cultivation.

References

- AOAC International (2005). Official Methods of Analysis of AOAC International. In: W. Horwitz (ed.), 18th edition, Arlington, Va.
- Basappa S.C., Shantha T. (1996). Methods for detoxification of aflatoxin in foods and feeds – a critical appraisal. *Journal of Food Science* 33(2): 95-107.
- Binder E.M. (2007). Managing the risk of mycotoxins in modern feed production. *Animal Feed Science and Technology* 133: 149-166.
- Brown R.L., Chen Z.Y., Cleveland T.E., Russin J.S. (1999). Advances in the development of host resistance in corn to aflatoxin contamination by *Aspergillus flavus*. *Phytopathology* 89(2): 113-117.
- Bullerman L.B., Liew F.Y., Seier S.A. (1977). Inhibition of growth and aflatoxin production by cinnamon and clove oils, cinnamic aldehyde and eugenol. *Journal of Food Science Chicago* 42(6): 1107-1109.
- CAST Report (2003). Mycotoxins, risk in plant, animal and human systems. In: J.L. Richard, G.A. Payne (eds.), Council of Agricultural Science and Technology Task Force, Report No. 139, Ames, Iowa, USA, ISBN 1-887383-22-0.
- Chen X.L., Zhou Y.D., Qu L.B., Liu X.Q., Yuan J.W., Zhao Y.F. (2010). Synthesis of Novel Coumarin-7,8-cyclophosphoramidate Analogs. *Synthetic Communications* 40(13): 1992-1997.
- Cho J., Kang G. (2000). Control of aflatoxin B₁ production of *Aspergillus parasiticus* using antagonistic microorganisms and its applications in Meju. *Food Science and Biotechnology* 9: 151-156.
- CODEX: CAC/RCP (51-2003, 2005). Code of practice for prevention and reduction of mycotoxin contamination in cereals. http://www.ipfsaph.org/eds_upload/kopool_data/codex_0/en_exc/051e.pdf.
- Delaš F., Duraković S., Sviben I.A., Martinko S., Markov K., Duraković L., Čvek D. (2008a). Antifungal Activity of Coumarin. In: D. Čurić (ed.). Proceedings of the Joint 4th Central European Congress on Food and 6th Croatian Congress of Food Technologists, Biotechnologists and Nutritionists, CEFood, Vol. 2., Cavtat, Croatia, pp 397-401.
- Delaš F., Duraković S., Sviben I.A., Martinko S., Markov K., Duraković L., Čvek D. (2008b). Antifungal Activity of Coumarin. Book of Abstracts of the Joint 4th Central European Congress on Food and 6th Croatian Congress of Food Technologists, Biotechnologists and Nutritionists, CEFood, Cavtat, Croatia, p 221.
- Delaš F., Markov K., Duraković L., Čvek D. (2008c). Inhibition of *Aspergillus flavus* culture growth and aflatoxin accumulation *in vitro* by coumarin derivative. Book of Abstracts of 14th World Congress of Food Science & Technology, Shanghai, China, p 63.
- Diener U.L., Cole R.J., Sanders T.H., Payne G.A., Lee L.S., Klich M.A. (1987). Epidemiology of aflatoxin formation by *Aspergillus flavus*. *Annual Review of Phytopathology* 25: 249-270.
- Donald W., Mirocha C.J. (1977). Chitin as a measure of fungal growth in stored corn and soybean seed. *Cereal Chemistry* 64: 466-474.
- Dorner J.W., Cole J.R., Wicklow D.T. (1999). Aflatoxin reduction in corn through field application of competitive fungi. *Journal of Food Protection* 62(6): 650-656.
- Duraković L., Petrović Z., Radić B., Duraković S. (1999). Antifungal and Antimycotoxigenic Properties of Newly Synthesized Derivatives of Tetraketone. *Kemija u Industriji* 48(4): 139-144.

- Duraković L., Delaš F., Markov K., Duraković S. (2004). Growth and Biosynthesis of Aflatoxin in the Presence of Newly Synthesized Analogues of Dehydroacetic Acid. In: D. Karlović (ed.). Proceedings of the 5th Croatian Congress of Food Technologists, Biotechnologists and Nutritionists, 2nd Central European Meeting, Opatija, Croatia, pp 366-370.
- Duraković L. (2007). The influence of chosen parameters on the growth of the mold *Aspergillus flavus* ATCC 26949 in mixed culture and biosynthesis of aflatoxins B₁ and G₁. Ph.D. thesis, Faculty of Food Technology and Biotechnology, University of Zagreb.
- Duraković L., Delaš F., Glancer-Šoljan M., Berković K., Čvek D., Markov K. (2007). Inhibition of aflatoxin accumulation by newly synthesized analogues of coumarine in *Aspergillus flavus* strains. Book of Abstracts of Central European Symposium of Industrial Microbiology and Microbial Ecology, Power of Microbes in Industry and Environment with International Participation, Zadar, Croatia, p 74.
- Duraković L., Delaš F., Blažinkov M., Šećerkadić L., Frece J., Redžepović S., Bošnjak M., Skelin A., Duraković Z., Duraković S. (2010). Aflatoxin Accumulation During the Growth of Mould *Aspergillus flavus* ATCC 26949 on Corn in Pure and Mixed Culture as Related to Various Temperature and Moisture Content. Croatian Journal of Food Technology, Biotechnology and Nutrition 5(1-2): 10-17.
- Duraković S. (1981). The influence of mixed mould cultures from the crops surface on biosynthesis of aflatoxins with mould *Aspergillus parasiticus* NRRL 2999. Ph.D. thesis, Technical Faculty, University of Zagreb.
- Duraković S., Duraković Z., Beritić T., Radić B., Lalić Lj., Delaš F. (1985). Biosynthesis of aflatoxins by *Aspergillus parasiticus* on roasted coffee beans. Periodicum Biologorum 87: 503-510.
- Duraković S., Sušnik I., Golem F.V., Duraković Z., Beritić T., Radić B., Filipović-Kovačević Ž., Pavlaković Z. (1994). Dehydroacetic Acid and the Newly Synthesized Schiff Base to Control Aflatoxin Accumulation. Kemija u Industriji 43(1): 7-12.
- Duraković S., Pavlaković Z., Filipović-Kovačević Ž., Duraković L., Šarac-Arneri R., Lalić Lj.M., Colić I. (1995). Antifungal and Antimycotoxigenic Properties of Newly Synthesized Derivatives of Tetraketone. In: G. Sontag (ed.). Proceedings of the Eight European Conference on Food Chemistry, EURO FOOD CHEM VIII, Vol. 2., Vienna, Austria, pp 320-323.
- Duraković S., Delaš F., Govori S., Kaljaj V., Rapić V., Kaljaj L., Halt M., Duraković L. (2006). Antifungal and antimycotoxigenic properties of newly synthesized derivatives of coumarine. Book of Abstracts of 2nd International Scientific Congress Food Sciences and Food Biotechnology in Developing Countries, Saltillo, Mexico, p 37.
- Duraković S., Duraković L., Vahčić N., Skelin A., Duraković Z. (2008). Affect of moulds growth in mixed cultures on production of aflatoxins on maize hybrid. Cereal Research Communications 36(Part 3, Supplement 5): 1615-1618.
- Georgopapadaku N.H. (1998). Antifungals: mechanism of action and resistance, established and novel drugs. Current Opinion in Microbiology 1: 547-557.
- Goldblatt L.A. (1969). Aflatoxin. Scientific Background, Control and Implications. L.A. Goldblatt (ed.), Academic Press, New York and London, 1-54.
- Gourama H., Bullerman L.B. (1995). *Aspergillus flavus* and *Aspergillus parasiticus*: Aflatoxigenic fungi of concern in foods and feeds: A review. Journal of Food Protection 59: 1395-1404.
- Govori S., Kaljaj V., Rapić V., Kaljaj L., Đaković S. (2002). Synthesis and structure of some 3,4-annelated coumarin systems. Heterocyclic Communications 8(2): 129-134.
- Hitokoto H., Murozumi S., Wauke T., Skai S., Ueno J. (1980). Inhibitory effects of condiments and herbal drugs on the growth and toxin production of toxigenic fungi. Mycopathologia 66: 167-177.
- Karlowy P. (1999). Biological detoxification of fungal toxins and its use in plant breeding, feed and food production. Natural Toxins 7(1): 1-23.
- Lavermicocca P., Valerio F., Visconti A. (2003). Antifungal activity of phenyllactic acid against molds isolated from bakery products. Applied and Environmental Microbiology 69(1): 634-640.
- Parameshwarappa G., Lingamani J., Patil S.B., Goudgaon N.M. (2009). Synthesis and antimicrobial activity of thiazole substituted coumarins. Heterocyclic Communications 15(5): 343-348.
- Pittet A. (1998). Natural occurrence of mycotoxins in foods and feeds. Revue de Médecine Vétérinaire 149(6): 479-492.
- Praxton J.D. (1991). Assays of antifungal activity. Methods in Plant Biochemistry 6: 33-46.
- Roussaki M., Kontoglorgis C.A., Hadjipaviou-Litina D., Hamilakis S., Detsi A. (2010). A novel synthesis of 3-aryl coumarins and evaluation of their antioxidant and lipoxigenase inhibitory activity. Bioorganic & Medicinal Chemistry Letters 20(13): 3889-3892.
- Samarajeva U., Seu A.C., Cohae M.D., Wie C.I. (1990). Detoxification of aflatoxins in foods and feeds by physical and chemical methods. Journal of Food Protection 53: 489-501.
- Schatzmayr G., Zehner F., Schatzmayr D., Täubel M., Binder E.M. (2006). Microbiologicals for deactivating mycotoxins in contaminated feed. Molecular Nutrition & Food Research 50(6): 543-551.
- Sušac K., Duraković S., Rapić V. (1989). Antifungal properties of newly synthesized derivatives of coumarine. Microbiologia 26(2): 165-172.
- Tamura S., Fujitani T., Kaneko M., Murakami N. (2010). Phenylcoumarin with Rev-export inhibitory activity from *Cnidii Monnieris Fructus*. Bioorganic & Medicinal Chemistry Letters 20(12): 3717-3720.
- Thompson C., Henke S.E. (2000). Effect of climate and type of storage container on aflatoxin production in corn and its associated risk to wildlife. Journal of Wildlife Diseases 36(1): 172-179.
- Weidenbörner M. (2001). Encyclopedia of Food Mycotoxins. Springer-Verlag, Berlin, ISBN 3540675566.
- Wicklow D.T., Weaver D.K., Throne J.E. (1998). Fungal colonists of maize grain conditioned at constant temperatures and humidities. Journal of Stored Products Research 34(4): 355-361.
- Wilkins K.M., Board R.G. (1991). Natural antimicrobial systems. In: Mechanisms of action of food preservation procedures. G.W. Gould (ed.), Elsevier, London, 285-362.
- Windham G.L., Williams W.P., Davis F.M. (1999). Effects of the southwestern corn borer on *Aspergillus flavus* kernel infection and aflatoxin accumulation in maize hybrids. Plant Disease 83(6): 535-540.
- Wright M.S., Greene-McDowelle D.M., Zeringue H.J., Bhatnagar D., Cleveland T.E. (2000). Effects of aldehydes from *Aspergillus*-resistant varieties of corn on *Aspergillus parasiticus* growth and aflatoxin biosynthesis. Toxicon 38(9): 1215-1223.
- Wu F. (2004). Mycotoxin risk assesment for the purpose of setting international regulatory standards. Environmental Science & Technology 38(15): 4049-4055.

acs75_24